

Filed on 23 August 2000

ADEOKUN, Miss Monisola,  
AMBROSE, Dr Helen Jean,  
CRESSWELL, Dr Carl John all British  
and all of AstraZeneca, Alderley Park,  
Macclesfield, Cheshire, GB-SK10 4TG  
and  
DUDLEY, Dr Adam Jeston,, Scientist  
(British)  
of AstraZeneca Pharmaceuticals,  
1800 Concord Pike, Wilmington,  
US-DE19850-5437;

## CHEMICAL COMPOUNDS

AstraZeneca AB,  
S-151 85 Södertälje, Sweden.

CHEMICAL COMPOUNDS

ins 171

This invention relates to polymorphisms in the human OATPC gene and corresponding novel allelic polypeptides encoded thereby. The invention also relates to methods and materials for analysing allelic variation in the OATPC gene, and to the use of OATPC polymorphism in treatment of diseases with OATPC transportable drugs.

Na<sup>+</sup>-independent organic anion transporting polypeptide (OATP) C gene is a member of the OATP supergene family involved in multifunctional transport of organic anion.

OATPC transports the organic anion taurocholate, conjugated steroids: DHEAS, estradiol 17 $\beta$ -

10 D-glucuronide and estrone-3-sulfate, eicosanoids: PGE<sub>2</sub>, thromboxane B<sub>2</sub>, leukotriene C<sub>4</sub>, and E<sub>4</sub>, and thyroid hormones T<sub>4</sub> and T<sub>3</sub><sup>1,2</sup>. OATPC has also been shown to be involved in the transport of xenobiotics, and drugs involved in lipid lowering *e.g.* statins<sup>1</sup>. Statins have been referred to as a first-line therapy for patients with atherosclerotic vascular diseases. The OATPC gene and its product is also thought to be of importance in other diseases due to its

15 transport of DHEAS an adrenal steroid which has been suggested to have positive neuropsychiatric, immune, and metabolic effects<sup>3</sup>. Due to the substrate specificity, location in the liver, and being exclusively expressed in the liver, Abe *et al* suggested that OATPC could be the predominant clearance mechanism of several endogenous and exogenous substrates in the liver. OATPC is the first human molecule reported to transport thyroid hormones<sup>2</sup>.

20 This liver specific transporter may be useful in liver-specific drug delivery systems and liver-specific chemotherapy, bile acid formation and the pathogenesis of diseases such as cholestasis, hyperbilirubinemia and thyroid hormone resistance.

The OATPC gene (sometimes called OAPT2 in the literature) has been cloned by four different groups, annotated and published as EMBL accession numbers AB026257 (OATPC, 25 2452bp), AF205071(OATP2, 2830, ref 1), AJ132573(OATP2, 2778)<sup>4</sup>, and AF060500 (LST-

<sup>1</sup> A Novel Human Hepatic Organic Anion Transporting Polypeptide (OATP2), Hsiang *et al* J Biol Chem **274**, 37161-37168 (1999)

<sup>2</sup> Identification of a Novel Gene Family Encoding Human Liver-specific Organic Anion Transporter LST-1, Takaaki Abe *et al* J Biol Chem **274**, 17159-17163 (1999)

<sup>3</sup> Bates *et al* (1998) Curr. Opin. Endocrinol. Diab. **5**, 357-366

<sup>4</sup> A novel human organic anion transporting polypeptide localised to the basolateral hepatocyte membrane, Konig Jorg *et al* (2000) Am J Physiol. Gastrointest. Liver Physiol. **278**: G156-G164

1)<sup>2</sup>. Polymorphism has been reported by Tamai<sup>5</sup> which is Asn130Asp and Val174Ala although any functional effect was stated therein to be not clear. König (2000) J Biol Chem 275, 23161-23168 describes the genomic organisation of OATP 1, 2 and 8. International patent application WO 00/08157 describes human anion transporter genes and some  
5 polymorphisms.

All positions herein of polymorphisms in the OATPC polynucleotide relate to the position in one of SEQ ID NO 1 or 3-12 unless stated otherwise or apparent from the context.

All positions herein of polymorphisms in the OATPC polypeptide relate to the position in SEQ ID NO 2 unless stated otherwise or apparent from the context.

10 One approach is to use knowledge of polymorphisms to help identify patients most suited to therapy with particular pharmaceutical agents (this is often termed "pharmacogenetics"). Pharmacogenetics can also be used in pharmaceutical research to assist the drug selection process. Polymorphisms are used in mapping the human genome and to elucidate the genetic component of diseases. The reader is directed to the following  
15 references for background details on pharmacogenetics and other uses of polymorphism detection: Linder *et al.* (1997), Clinical Chemistry, **43**, 254; Marshall (1997), Nature Biotechnology, **15**, 1249; International Patent Application WO 97/40462, Spectra Biomedical; and Schafer *et al.* (1998), Nature Biotechnology, **16**, 33.

Clinical trials have shown that patient response to treatment with pharmaceuticals is  
20 often heterogeneous. Thus there is a need for improved approaches to pharmaceutical agent design and therapy.

Point mutations in polypeptides will be referred to as follows: natural amino acid (using 1 or 3 letter nomenclature), position, new amino acid. For (a hypothetical) example "D25K" or "Asp25Lys" means that at position 25 an aspartic acid (D) has been changed to  
25 lysine (K). Multiple mutations in one polypeptide will be shown between square brackets with individual mutations separated by commas.

The present invention is based on the discovery of polymorphisms in OATPC.

According to one aspect of the present invention there is provided a method for the detection of a polymorphism in OATPC in a human, which method comprises determining the  
30 sequence of the human at at least one polymorphic position and determining the status of the

<sup>5</sup> Tamai *et al.* (2000), BBRC, **273**, 251-60

A method for the detection of a polymorphism in OATPC in a human, which method comprises determining the sequence of the human at at least one of the following polymorphic

positions 510, 696, 1299, 1312, 1347, 1561, 2028, 2327 and 2342 in sequence of the OATPC gene as defined by the position in SEQ ID NO: 1;

positions 400, 405, 488 and 643 in OATPC polypeptide defined by position in SEQ ID NO: 2;

positions 321 and 1332 defined by position in SEQ ID NO 3;

10 position 41 defined by position in SEQ ID NO 4;

positions 109 and 244 defined by position in SEQ ID NO 5;

positions 117 and 283 defined by position in SEQ ID NO 6;

positions 209 and 211 defined by position in SEQ ID NO 7;

positions 63 to 68 defined by position in SEQ ID NO 8;

15 position 53 defined by position in SEQ ID NO 9;

position 75 defined by position in SEQ ID NO 10;

position 162 defined by position in SEQ ID NO 11; and

positions 84 defined by position in SEQ ID NO 12.

The term human includes both a human having or suspected of having a OATPC mediated disease and an asymptomatic human who may be tested for predisposition or susceptibility to such disease. At each position the human may be homozygous for an allele or the human may be a heterozygote.

The term polymorphism includes single nucleotide substitution, nucleotide insertion and nucleotide deletion which in the case of insertion and deletion includes insertion or deletion of one or more nucleotides at a position of a gene.

The method for diagnosis is preferably one in which the sequence is determined by a method selected from amplification refractory mutation system and restriction fragment length polymorphism.

The status of the individual may be determined by reference to allelic variation at any 30 one, two, three, four, five, six, seven, eight, nine or more positions.

The test sample of nucleic acid is conveniently a sample of blood, bronchoalveolar lavage fluid, sputum, or other body fluid or tissue obtained from an individual. It will be

appreciated that the test sample may equally be a nucleic acid sequence corresponding to the sequence in the test sample, that is to say that all or a part of the region in the sample nucleic acid may firstly be amplified using any convenient technique e.g. PCR, before analysis of allelic variation.

- 5 It will be apparent to the person skilled in the art that there are a large number of analytical procedures which may be used to detect the presence or absence of variant nucleotides at one or more polymorphic positions of the invention. In general, the detection of allelic variation requires a mutation discrimination technique, optionally an amplification reaction and optionally a signal generation system. Table 1 lists a number of mutation
- 10 detection techniques, some based on the PCR. These may be used in combination with a number of signal generation systems, a selection of which is listed in Table 2. Further amplification techniques are listed in Table 3. Many current methods for the detection of allelic variation are reviewed by Nollau *et al.*, Clin. Chem. 43, 1114-1120, 1997; and in standard textbooks, for example "Laboratory Protocols for Mutation Detection", Ed. by U.
- 15 Landegren, Oxford University Press, 1996 and "PCR", 2<sup>nd</sup> Edition by Newton & Graham, BIOS Scientific Publishers Limited, 1997.

#### Abbreviations:

ALEX™	Amplification refractory mutation system linear extension
APEX	Arrayed primer extension
ARMST™	Amplification refractory mutation system
b-DNA	Branched DNA
bp	base pair
CMC	Chemical mismatch cleavage
COPS	Competitive oligonucleotide priming system
DGGE	Denaturing gradient gel electrophoresis
FRET	Fluorescence resonance energy transfer
HMG-CoA	3-hydroxy-3-methylglutaryl-coenzyme A
LCR	Ligase chain reaction
MASDA	Multiple allele specific diagnostic assay
NASBA	Nucleic acid sequence based amplification
OATP	Na <sup>+</sup> -independent organic anion transporting polypeptide
OLA	Oligonucleotide ligation assay
PCR	Polymerase chain reaction
PTT	Protein truncation test
RFLP	Restriction fragment length polymorphism
SDA	Strand displacement amplification
SNP	Single nucleotide polymorphism
SSCP	Single-strand conformation polymorphism analysis

SSR	Self sustained replication
TGGE	Temperature gradient gel electrophoresis

### Table 1 - Mutation Detection Techniques

**General:** DNA sequencing, Sequencing by hybridisation

**Scanning:** PTT\*, SSCP, DGGE, TGGE, Cleavase, Heteroduplex analysis, CMC, Enzymatic mismatch cleavage

\* Note: not useful for detection of promoter polymorphisms.

#### **Hybridisation Based**

Solid phase hybridisation: Dot blots, MASDA, Reverse dot blots,  
Oligonucleotide arrays (DNA Chips)

10 Solution phase hybridisation: Taqman™ - US-5210015 & US-5487972 (Hoffmann-La Roche), Molecular Beacons - Tyagi *et al* (1996), Nature Biotechnology, **14**, 303; WO 95/13399 (Public Health Inst., New York)

**Extension Based:** ARMS™, ALEX™ - European Patent No. EP 332435 B1 (Zeneca Limited), COPS - Gibbs *et al* (1989), Nucleic Acids Research, **17**, 2347.

15 **Incorporation Based:** Mini-sequencing, APEX

**Restriction Enzyme Based:** RFLP, Restriction site generating PCR

**Ligation Based:** OLA

**Other:** Invader assay

### 20 Table 2 - Signal Generation or Detection Systems

**Fluorescence:** FRET, Fluorescence quenching, Fluorescence polarisation - United Kingdom Patent No. 2228998 (Zeneca Limited)

**Other:** Chemiluminescence, Electrochemiluminescence, Raman, Radioactivity, Colorimetric, Hybridisation protection assay, Mass spectrometry

25

### Table 3 - Further Amplification Methods

SSR, NASBA, LCR, SDA, b-DNA

Preferred mutation detection techniques include ARMS™, ALEX™, COPS, Taqman,

30 Molecular Beacons, RFLP, and restriction site based PCR and FRET techniques.

Particularly preferred methods include ARMS™ and RFLP based methods. ARMS™ is an especially preferred method.

In a further aspect, the diagnostic methods of the invention are used to assess the pharmacogenetics of a drug transportable by OATPC.

- 5 Assays, for example reporter-based assays, may be devised to detect whether one or more of the above polymorphisms affect transcription levels and/or message stability.

Individuals who carry particular allelic variants of the OATPC gene may therefore exhibit differences in their ability to regulate protein biosynthesis under different physiological conditions and will display altered abilities to react to different diseases. In addition, differences arising as a result of allelic variation may have a direct effect on the response of an individual to drug therapy. The diagnostic methods of the invention may be useful both to predict the clinical response to such agents and to determine therapeutic dose.

- 10 In a further aspect, the diagnostic methods of the invention, are used to assess the predisposition and/or susceptibility of an individual to diseases mediated by OATPC. This may be particularly relevant in the development of hyperlipoproteinemia and cardiovascular disease and the present invention may be used to recognise individuals who are particularly at risk from developing these conditions.

In a further aspect, the diagnostic methods of the invention are used in the development of new drug therapies which selectively target one or more allelic variants of the OATPC gene. Identification of a link between a particular allelic variant and predisposition to disease development or response to drug therapy may have a significant impact on the design of new drugs. Drugs may be designed to regulate the biological activity of variants implicated in the disease process whilst minimising effects on other variants.

- 20 In a further diagnostic aspect of the invention the presence or absence of variant nucleotides is detected by reference to the loss or gain of, optionally engineered, sites recognised by restriction enzymes.

According to another aspect of the present invention there is provided a human OATPC gene or its complementary strand comprising a variant allelic polymorphism at one or more of positions defined herein or a fragment thereof of at least 20 bases comprising at least one novel polymorphism.

30 Fragments are at least 17 bases, more preferably at least 20 bases, more preferably at least 30 bases.

According to another aspect of the present invention there is provided a polynucleotide comprising at least 20 bases of the human OATPC gene and comprising an allelic variant selected from any one of the following:

Region	variant	Position in SEQ ID NO	SEQ ID NO
Exon 4	A	510	1
Exon 5	T	670	1
Exon 5	T	696	1
Exon 9	G	1299	1
Exon 9	A	1312	1
Exon 9	A	1347	1
Exon 10	C	1561	1
Exon 14	C	2028	1
3'UTR	Insert T	2327	1
3'UTR	C	2342	1
Promoter	G	321	3
Promoter	C	1332	3
Intron 1	A	41	4
Intron 2	G	109	5
Intron 2	G	244	5
Intron 3	A	117	6
Intron 3	A	283	6
Intron 4	A	209	7
Intron 4	A	211	7
Intron 4	Deletion CTTGTA	63	8
Intron 6	T	53	9
Intron 9	Insert TTC	75	10
Intron 11	Insert T	162	11
Intron 12	C	84	12

- 5 According to another aspect of the present invention there is provided a human OATPC gene or its complementary strand comprising a polymorphism, preferably corresponding with one or more the positions defined herein or a fragment thereof of at least 20 bases comprising at least one polymorphism.

Fragments are at least 17 bases, more preferably at least 20 bases, more preferably at least 30 bases.

The invention further provides a nucleotide primer which can detect a polymorphism of the invention.



According to another aspect of the present invention there is provided an allele specific primer capable of detecting a OATPC gene polymorphism, preferably at one or more of the positions as defined herein.

An allele specific primer is used, generally together with a constant primer, in an amplification reaction such as a PCR reaction, which provides the discrimination between alleles through selective amplification of one allele at a particular sequence position e.g. as used for ARMS™ assays. The allele specific primer is preferably 17- 50 nucleotides, more preferably about 17-35 nucleotides, more preferably about 17-30 nucleotides.

An allele specific primer preferably corresponds exactly with the allele to be detected but derivatives thereof are also contemplated wherein about 6-8 of the nucleotides at the 3' terminus correspond with the allele to be detected and wherein up to 10, such as up to 8, 6, 4, 2, or 1 of the remaining nucleotides may be varied without significantly affecting the properties of the primer.

Primers may be manufactured using any convenient method of synthesis. Examples of such methods may be found in standard textbooks, for example "Protocols for Oligonucleotides and Analogues; Synthesis and Properties," Methods in Molecular Biology Series; Volume 20; Ed. Sudhir Agrawal, Humana ISBN: 0-89603-247-7; 1993; 1<sup>st</sup> Edition. If required the primer(s) may be labelled to facilitate detection.

According to another aspect of the present invention there is provided an allele-specific oligonucleotide probe capable of detecting a OATPC gene polymorphism, preferably at one or more of the positions defined herein.

The allele-specific oligonucleotide probe is preferably 17- 50 nucleotides, more preferably about 17-35 nucleotides, more preferably about 17-30 nucleotides.

The design of such probes will be apparent to the molecular biologist of ordinary skill. Such probes are of any convenient length such as up to 50 bases, up to 40 bases, more conveniently up to 30 bases in length, such as for example 8-25 or 8-15 bases in length. In general such probes will comprise base sequences entirely complementary to the corresponding wild type or variant locus in the gene. However, if required one or more mismatches may be introduced, provided that the discriminatory power of the oligonucleotide probe is not unduly affected. The probes of the invention may carry one or more labels to facilitate detection.

According to another aspect of the present invention there is provided an allele specific primer or an allele specific oligonucleotide probe capable of detecting a OATPC gene polymorphism at one of the positions defined herein.

According to another aspect of the present invention there is provided a diagnostic kit  
5 comprising an allele specific oligonucleotide probe of the invention and/or an allele-specific primer of the invention.

The diagnostic kits may comprise appropriate packaging and instructions for use in the methods of the invention. Such kits may further comprise appropriate buffer(s) and polymerase(s) such as thermostable polymerases, for example taq polymerase.

10 In another aspect of the invention, the single nucleotide polymorphisms of this invention may be used as genetic markers in linkage studies. This particularly applies to the polymorphisms of relatively high frequency. The OATPC gene is on chromosome 12p (as shown from a database search with the cDNA as a query sequence). Low frequency polymorphisms may be particularly useful for haplotyping as described below. A haplotype is  
15 a set of alleles found at linked polymorphic sites (such as within a gene) on a single (paternal or maternal) chromosome. If recombination within the gene is random, there may be as many as  $2^n$  haplotypes, where 2 is the number of alleles at each SNP and n is the number of SNPs. One approach to identifying mutations or polymorphisms which are correlated with clinical response is to carry out an association study using all the haplotypes that can be identified in  
20 the population of interest. The frequency of each haplotype is limited by the frequency of its rarest allele, so that SNPs with low frequency alleles are particularly useful as markers of low frequency haplotypes. As particular mutations or polymorphisms associated with certain clinical features, such as adverse or abnormal events, are likely to be of low frequency within the population, low frequency SNPs may be particularly useful in identifying these mutations  
25 (for examples see: Linkage disequilibrium at the cystathionine beta synthase (CBS) locus and the association between genetic variation at the CBS locus and plasma levels of homocysteine. *Ann Hum Genet* (1998) 62:481-90, De Stefano V, Dekou V, Nicaud V, Chasse JF, London J, Stansbie D, Humphries SE, and Gudnason V; and Variation at the von willebrand factor (vWF) gene locus is associated with plasma vWF:Ag levels: identification of three novel  
30 single nucleotide polymorphisms in the vWF gene promoter. *Blood* (1999) 93:4277-83, Keightley AM, Lam YM, Brady JN, Cameron CL, Lillicrap D).

According to another aspect of the present invention there is provided a computer readable medium comprising at least one novel sequence of the invention stored on the medium. The computer readable medium may be used, for example, in homology searching, mapping, haplotyping, genotyping or pharmacogenetic analysis.

5 According to another aspect of the present invention there is provided a method of treating a human in need of treatment with a drug transportable by OATPC in which the method comprises:

i) detection of a polymorphism in OATPC in the human, which detection comprises determining the sequence of the human at one or more of the following positions:

10 positions 487, 510, , 554, 670, 696, 819, 820, 1299, 1312, 1347, 1561, 2028, 2327 and 2342 in sequence of the OATPC gene as defined by the position in SEQ ID NO: 1;

positions 130, 152, 174, 241, 400, 405, 488 and 643 in OATPC polypeptide defined by position in SEQ ID NO: 2;

positions 321 and 1332 defined by position in SEQ ID NO 3;

15 position 41 defined by position in SEQ ID NO 4;

positions 109 and 244 defined by position in SEQ ID NO 5;

positions 117 and 283 defined by position in SEQ ID NO 6;

positions 209 and 211 defined by position in SEQ ID NO 7;

positions 63 to 68 defined by position in SEQ ID NO 8;

20 position 53 defined by position in SEQ ID NO 9;

position 75 defined by position in SEQ ID NO 10;

position 162 defined by position in SEQ ID NO 11; and

positions 84 defined by position in SEQ ID NO 12.

and determining the status of the human by reference to polymorphism in the OATPC gene;

25 and

ii) administering an effective amount of the drug.

Preferably determination of the status of the human is clinically useful. Examples of clinical usefulness include deciding which statin drug or drugs to administer and/or in deciding on the effective amount of the statin drug or drugs. Statins already approved for use  
30 in humans include atorvastatin, cerivastatin, fluvastatin, pravastatin and simvastatin. The reader is referred to the following references for further information: Drugs and Therapy Perspectives (12<sup>th</sup> May 1997), 9: 1-6; Chong (1997) Pharmacotherapy 17: 1157-1177; Kellick

- (1997) Formulary **32**: 352; Kathawala (1991) Medicinal Research Reviews, **11**: 121-146; Jahng (1995) Drugs of the Future **20**: 387-404, and Current Opinion in Lipidology, (1997), **8**, 362 - 368. A preferred statin drug is compound 3a (S-4522) in Watanabe (1997) Bioorganic and Medicinal Chemistry **5**: 437-444; now called rosuvastatin, see Olsson (2001) American Journal of Cardiology, **87**, supplement 1, 33-36. The term "drug transportable by OATPC" means that transport by OATPC in humans is an important part of a drug exerting its pharmaceutical effect in man. For example, some statins have to be transported to the liver by OATPC to exert their lipid lowering effects.

- According to another aspect of the present invention there is provided a method of treating a human in need of treatment with a drug transportable by OATPC in which the method comprises:
- i) diagnosis of a single nucleotide polymorphism in OATPC gene in the human, which diagnosis preferably comprises determining the sequence of the nucleic acid at position 487 in the coding sequence of the OATPC gene as defined by the position in SEQ ID NO: 1.
  - and determining the status of the human by reference to polymorphism in the OATPC gene; and
  - ii) administering an effective amount of the drug.

- Although the polymorphism at position 487 (A→G; Asn130Asp) was reported by Tamai, no functional effect was attributed. Indeed the following polymorphisms are known but have not previously been attributed a role in statin pharmacogenetics as disclosed herein.

#### OATPC polymorphisms

Position of nucleotide as in sequence 1 of OATPC patent

Position of amino acid as in sequence 2 of OATPC patent

Exon	nucleotide	SNP	Aminoacid	Change	Change
4	487	A>G	130	Asn>Asp	N>D
4	554	G>A	152	Arg>Lys	R>K
5	620	T>C	174	Val>Ala	V>A
5	670	C>T	191	Leu	L
6	819	A>T	240	Val	V
6	820	G>A	241	Asp>Asn	D>N

According to another aspect of the present invention there is provided use of a drug transportable by OATPC in preparation of a medicament for treating a disease in a human

diagnosed as having a polymorphism therein, preferably at one or more of the positions defined herein. Preferably the disease is cardiovascular.

According to another aspect of the present invention there is provided a pharmaceutical pack comprising OATPC transportable drug and instructions for administration of the drug to humans diagnostically tested for a single nucleotide polymorphism therein, preferably at one or more of the positions defined herein.

According to another aspect of the present invention there is provided an allelic variant of human OATPC polypeptide comprising at least one of the following:

- a leucine at position 400 of SEQ ID NO 2;
- 10 an isoleucine at position 405 of SEQ ID NO 2;
- an arginine at position 488 of SEQ ID NO 2;
- a phenylalanine at position 643 of SEQ ID NO 2;
- or a fragment thereof comprising at least 10 amino acids provided that the fragment comprises at least one allelic variant.

- 15 Fragments of polypeptide are at least 10 amino acids, more preferably at least 15 amino acids, more preferably at least 20 amino acids.

According to another aspect of the present invention there is provided an antibody specific for an allelic variant of human OATPC polypeptide as described herein.

- Antibodies can be prepared using any suitable method. For example, purified
- 20 polypeptide may be utilized to prepare specific antibodies. The term "antibodies" is meant to include polyclonal antibodies, monoclonal antibodies, and the various types of antibody constructs such as for example  $F(ab')_2$ , Fab and single chain Fv. Antibodies are defined to be specifically binding if they bind the allelic variant of OATPC with a  $K_a$  of greater than or equal to about  $10^7 M^{-1}$ . Affinity of binding can be determined using conventional techniques,
  - 25 for example those described by Scatchard et al., *Ann. N.Y. Acad. Sci.*, 51:660 (1949).

- Polyclonal antibodies can be readily generated from a variety of sources, for example, horses, cows, goats, sheep, dogs, chickens, rabbits, mice or rats, using procedures that are well-known in the art. In general, antigen is administered to the host animal typically through parenteral injection. The immunogenicity of antigen may be enhanced through the use of an
- 30 adjuvant, for example, Freund's complete or incomplete adjuvant. Following booster immunizations, small samples of serum are collected and tested for reactivity to antigen. Examples of various assays useful for such determination include those described in:

*Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988; as well as procedures such as countercurrent immuno-electrophoresis (CIEP), radioimmunoassay, radioimmunoprecipitation, enzyme-linked immuno-sorbent assays (ELISA), dot blot assays, and sandwich assays, see U.S. Patent Nos. 4,376,110 and 4,486,530.

- 5 Monoclonal antibodies may be readily prepared using well-known procedures, see for example, the procedures described in U.S. Patent Nos. RE 32,011, 4,902,614, 4,543,439 and 4,411,993; *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, Kennett, McKearn, and Bechtol (eds.), (1980).

- The monoclonal antibodies of the invention can be produced using alternative  
10 techniques, such as those described by Altling-Mees et al., "Monoclonal Antibody Expression Libraries: A Rapid Alternative to Hybridomas", *Strategies in Molecular Biology* 3: 1-9 (1990) which is incorporated herein by reference. Similarly, binding partners can be constructed using recombinant DNA techniques to incorporate the variable regions of a gene that encodes a specific binding antibody. Such a technique is described in Larrick et al., *Biotechnology*, 7:  
15 394 (1989).

Once isolated and purified, the antibodies may be used to detect the presence of antigen in a sample using established assay protocols, see for example "A Practical Guide to ELISA" by D. M. Kemeny, Pergamon Press, Oxford, England.

- According to another aspect of the invention there is provided a diagnostic kit  
20 comprising an antibody of the invention.

The invention will now be illustrated but not limited by reference to the following Examples. All temperatures are in degrees Celsius.

In the Examples below, unless otherwise stated, the following methodology and materials have been applied.

- 25 AMPLITAQ™, available from Perkin-Elmer Cetus, is used as the source of thermostable DNA polymerase.

General molecular biology procedures can be followed from any of the methods described in "Molecular Cloning - A Laboratory Manual" Second Edition, Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory, 1989).

- 30 Electropherograms were obtained in a standard manner: data was collected by ABI377 data collection software and the wave form generated by ABI Prism sequencing analysis (2.1.2).

Example 1**Identification of Polymorphisms****1. Methods**DNA Preparation

- 5 DNA was prepared from frozen blood samples collected in EDTA following protocol I (Molecular Cloning: A Laboratory Manual, p392, Sambrook, Fritsch and Maniatis, 2<sup>nd</sup> Edition, Cold Spring Harbor Press, 1989) with the following modifications. The thawed blood was diluted in an equal volume of standard saline citrate instead of phosphate buffered saline to remove lysed red blood cells. Samples were extracted with phenol, then
- 10 phenol/chloroform and then chloroform rather than with three phenol extractions. The DNA was dissolved in deionised water.

Template Preparation

- Templates were prepared by PCR using the oligonucleotide primers and annealing temperatures set out below. The extension temperature was 72° and denaturation temperature
- 15 94°. Generally 50 ng of genomic DNA was used in each reaction and subjected to 35 cycles of PCR. Where described below, the primary fragment was diluted 1/100 and two microlitres were used as template for amplification of secondary fragments. PCR was performed in two stages (primary fragment then secondary fragment) to ensure specific amplification of the desired target sequence.
- 20 **Polymorphisms in OATPC: cDNA screening of 15 Liver samples**

Region	SNP	Position	Amino Acid Change	Allele frequencies
Exon 4	G/A	510	None	G=96.7% A=3.3%
Exon 5	C/T	670	None	C=50% T=50%
Exon 5	C/T	696	None	C=60% T=40%
Exon 9	C/G	1299	Phe400Leu	C=96.7% G=3.3%
Exon 9	G/A	1312	Val405Ile	G=96.7% A=3.3%
Exon 9	G/A	1347	None	G=96.7% A=3.3%
Exon 10	G/C	1561	Gly488Arg	G=96.7% C=3.3%
Exon 14	A/C	2028	Leu643Phe	A=90%

				C=10%
--	--	--	--	-------

OATP2 above refers to the clone sequenced by Hsiang et al (ref 1). Some comment on the numbering of exons in OATPC is required. This gene contains an exon (38 bp) upstream (5' UTR region) of the exon containing the ATG start site for translation. Therefore the exon numbering could vary depending whether this exon is counted as the first exon or not. In the literature, Konig (2000) JBC 275: 23161-68, have defined exon 1 as that containing the ATG start site and therefore we have adopted the same numbering in this application (but note that the priority document relating to the present application did vice versa; for example, exon 5 in this application is equivalent to exon 6 in the priority document).

## PCR PRODUCTS

Fragment	Forward Oligo	Reverse Oligo
443-999	443-466	979-999
874-1360	874-896	1337-1360
1255-1684	1255-1278	1663-1684
1559-2095	1559-1581	2073-2095

## 20 RFLP analysis

Polymorphism	Position	RFLP Enzyme/ PCR size	RFLP fragment size
G/A	510		
C/T	670	BmR I/ 595bp	C=349bp, 246bp T=595bp
C/T	696		
C/G	1299	Apo I/ 595bp	C=30bp, 60bp, 380bp G=90bp, 380bp
G/A	1312	Bst 4Cl/ 595bp	G=77bp, 393bp A=470bp
G/A	1347		
G/C	1561	HpyCH4IV/ 595bp	G=470bp C=144bp, 326bp
A/C	2028	Ase I/ 577bp	A=89bp, 488bp C=577bp



Example 2**Further OATPC Polymorphisms****SNPs in OATPC 3'UTR (positions according to SEQ ID NO 1)**

Exon	Nucleotide	SNP				Frequency Caucasian	Frequency Japanese
3' UTR	2327	Ins T				not screened	0.1
3' UTR	2342	T>C				not screened	0.4

**SNPs in OATPC promoter (positions according to SEQ ID NO 3)**

	Nucleotide	SNP				Frequency Caucasian	Frequency Japanese
	321	T>G				0.03	not screened
	1332	A>C				0.08	not screened

**SNPs in OATPC introns**

Intron	Nucleotide position in relation to exon	SNP	Nucleotide position in sequence	Sequence ID No		
1	IVS1+21	T>A	41	4		
2	IVS2+89	T>G	109	5		
2	IVS2+224	A>G	244	5		
3	IVS3+97	C>A	117	6		
3	IVS3+263	G>A	283	6		
4	IVS4+189	G>A	209	7		
4	IVS4+191	G>A	211	7		
4	IVS5-118	delCTTGTA	63	8		
6	IVS6+33	C>T	53	9		
9	IVS10-107	ins TTC	75	10		
11	IVS11+142	Ins T	162	11		
12	IVS13-97	G>C	84	12		

**5 OATPC intronic SNPs****Key**

20bp of exon sequence shown in uppercase

Intron sequence in lowercase (200 to 300bp only)

10 SNP shown in uppercase (one allele only)

Sequence ID No 4

IVS1+21 T&gt;A

SNP at position 41 in this sequence

15

GATACTGCAA TGGATTGAAG gtagaataag ttttatgttt Ttgagctaaa ataagtaaat 60

5

SNP at position 109 in this sequence

15

SNP at position 117 in this sequence

25

SNP at position 211 in this sequence

00

50

ggtcatcat caaaagcaaat ttctttcattat aaagc

10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044

Sequence ID No 12  
IVS13-97 G>C

SNP at position 84 in this sequence

5 tgaatttgggt ctttgagatt tctaataatc ttattatttg ggtagatgca gaacaaaata 60  
ataaacgaat cctccaaaat tttGaacttt tatttaatac aaatatatca atgtggaata 120  
tcattgcagtt acatttaaaa tatgttcctt aaactgacat ctctctctct cctattacag 180  
10 GAGGAATTCT AGCTCCAATA 200

### OATPC promoter region

Total length of the sequence = 1538bp

1500bp of OATPC sequence directly upstream from the cDNA sequence

Sequence in uppercase represents 38bp overlap with the cDNA sequence (SEQ ID NO 1)

15 where this 38bp is 5'UTR sequence.

Nucleotide positions in the promoter have been determined where the -1 position is the base (lowercase) directly upstream of the end of the cDNA sequence.

20 atgtctctttg acctctgaaa atattggaga attttacaac tggcaccttt agctcaggat 60  
tataaaaggtt gttagttagt ttgtactgtt ttactctcat tgcataaant atatatatta 120  
gtctccaaaac atgttgatgt gttttcaatg aaatggatgt ctgaggagaa aacattatgc 180  
ctgagaaaaac caaaactgta ttcccatgtt gaataaaagg aagtcataaa aaatgatgga 240  
aaatgtttctg cattcctgtt atgatataca aatctggcag tacatgaaaa tttttcaagg 300  
tgcttatttta acaggcataa tctttgggtc cctgagccag aatctgctgg gtatgggact 360  
25 ggattgtctat ttgacaact cgcacagtaga ttcttactca gcacagtatt tggagagcct 420  
actctaatat ttgggccttg ggtctacatt tctcagttct gcacagtcac tctcccccct 480  
tacactactc tttagtttgt cctcatgttc caatactctc aataattaac caagataaga 540  
actaatcaat cagataactg tggcacagac atcaataaca ttttcttgca accatatcaa 600  
caaatgtccc atgaatgata aggggttaacc atattctcat atatgcattc tcacattacc 660  
30 acataatgat atgtgataac gtgtatacag gtaaaagtgt gtatatatgt atacattgat 720  
gtttgttgtgt atatacatc atatacttc acactcttct gaatatatata tattttatgt 780  
agagaaggggt ctgtacttta ttccaagaaga gagcttaagt tccaaggatg aattgagagt 840  
ctaaaagtgt ttgagtattg aattaattaa acttcacttc tactcaagaa aacttttaac 900  
tgagttaagc tctctccctc tcccaagctc aagtcataaa aaggaaactg tgatattaat 960  
35 aattctttcc ttgttttgatg taaagaatct atcgcataaa gcagttctaa ttttcatcat 1020  
tcagaaaaat ggtcttgacg ttaattggga ctctcttatt ccaggtggta tctccagctc 1080  
ccatacatca caggttagaa ccatacttat gtaccaagca aagagggtat attttaattt 1140  
ttaatgccca atgtacaactg taggcattat ttattattgt cttaaatat tctctatttt 1200  
gaagttttaa atactcggaa taattttatt tactcatatt tttaaagaaa aaaaacttat 1260  
40 gccaccactc taatttgata aacaagtaaa agccattccc aaaaagtaag ttactcttgt 1320  
aagatttaaca aaaaataatg tgagaattctt gagaaaataa atctttaaat atttgcaact 1380  
ggagtgaact cttaaaacta actaggtttt atabgtttga ctgagacat gacataataa 1440  
ggtggttaac catcactgga cttgttttca aaaagccaac tacttttaaga ggaataaagg 1500  
45 GTGACTTGT TCGAGTGTCT GTAGATTCT AAATCCAG 1538